

Table: Social Class (Percentage of the Kingston Population, 2011 Census)

	Kingston	London	England
1. Higher managerial, administrative and professional occupations	16.1	13.2	10.4
2. Lower managerial, administrative and professional occupations	25.6	23.1	20.9
3. Intermediate occupations	12.3	11.8	12.8
4. Small employers and own account workers	9.4	9.4	9.4
5. Lower supervisory and technical occupations	4.2	5	6.9
6. Semi-routine occupations	9	10.4	14
7. Routine occupations	5.2	7.4	11
8. Never worked and long-term unemployed	4.5	8.3	5.6
L15 Full-time students	13.7	11.4	9

Source: 2011 Census, Office of National Statistics

Table: Air Pollution (Annual concentration of human-made fine particulate matter at an area level Mean - µg/m3, 2016)

	Kingston	London	England
Annual concentration of human-made fine particulate matter at an area level Mean	10.8	11.4	9.3

Source: Public Health England, 2018

Table: Fraction of all-cause adult mortality (persons aged 30 years and above) attributable to anthropogenic particulate air pollution (measured as fine particulate matter, PM2.5) in Kingston, London and England, 2010 to 2015

Period	Kingston	London	England
2011	6.7	7.2	5.4
2012	6.1	6.6	5.1
2013	6.2	6.7	5.3
2014	5.9	6.5	5.1
2015	5.1	5.6	4.7
2016	6.1	6.4	5.3

Note: PM_{2.5} means the mass (in micrograms) per cubic metre of air of individual particles with an aerodynamic diameter generally less than 2.5 micrometers. PM_{2.5} is also known as fine particulate matter

Note, attributable deaths statistic is an estimate of the overall health effect of air pollution on the population of an area. It is derived from the rise in risk of death associated with a standard level of pollution, then converted to an equivalent number of deaths based on population and pollution levels to help describe the scale of this risk.

Source: Public Health England, 2017

Table: Lone Parent Household (All lone parent households with dependent children under 16 where the lone parent is aged 16 to 74), 2011

	Kingston	London	England
Total %	5.6	8.5	7.1
Male	340	20,953	151,744
Female	3,201	256,636	1,412,937

Source: 2011 Census, Office of National Statistics

Table: Provision of unpaid care by number of hours and age in Kingston, 2011

Age	Provides unpaid care: Total		Provides 1 to 19 hours unpaid care a week		Provides 20 to 49 hours unpaid care a week		Provides 50 or more hours unpaid care a week	
	Count	%	Count	%	Count	%	Count	%
All Ages	13,288	8.30%	9,331	5.80%	1,611	1.00%	2,346	1.50%
Age 0 to 15	251	0.80%	206	0.70%	19	0.10%	26	0.10%
Age 16 to 24	786	3.60%	629	2.90%	108	0.50%	49	0.20%
Age 25 to 34	1,229	4.80%	852	3.30%	184	0.70%	193	0.80%
Age 35 to 49	3,705	10.10%	2,569	7.00%	459	1.30%	677	1.80%
Age 50 to 64	4,787	18.80%	3,630	14.30%	548	2.20%	609	2.40%
Age 65 and over	2,530	12.40%	1,445	7.10%	293	1.40%	792	3.90%

Source: 2011 Census, Office of National Statistics

Table: Percentage of Unpaid Carers and Social Isolation among Adult Carers and All Carers

	Period	Kingston	London	England	
Unpaid Carers (% of population)	18+	2011	1.47	1.83	2.37
Social isolation among carers*	18+	2016/17	27.8	35.6	35.5

Note: Percentage of adult carers who have as much social contact at they would like according to the Personal Social Services Carers survey

Source: 2011 Census, Office of National Statistics & Public Health England, 2018

Table: Reported Quality of Life, Overall satisfaction with Social Services, and Adult Carers receiving Assessments, Kingston, London, and England

	Kingston	London	England
Carer reported quality of life (%) 2014/15	7.5	7.6	7.9
Overall satisfaction of carers with social services (%) 2014/15	32.2	35.2	41.2

Adult carers receiving assessments / 100,000 (2013/14)	252	684	968
--	-----	-----	-----

Source: Public Health England, 2018

Table: Percentage of eligible Adults with a Learning Disability who have had a GP Annual Health Checks (2016/17)

Kingston	London	England
51.40%	48.40%	48.90%
(45.6 to 57.7)	(47.4 to 49.2)	(48.6 to 49.1)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England, 2018

Table: Projections for Numbers of Adults aged 18-64 in Kingston predicted to have Autistic Spectrum Disorder

	2017	2020	2025	2030	2035
People aged 18-24 predicted to have ASD	177	179	185	211	219
People aged 25-34 predicted to have ASD	288	289	288	280	292
People aged 35-44 predicted to have ASD	288	298	312	317	314
People aged 45-54 predicted to have ASD	238	252	268	284	294
People aged 55-64 predicted to have ASD	172	186	208	224	238
Total population aged 18-64 predicted to have ASD	1163	1204	1261	1317	1358

Source: Projecting Older People Population Information (www.POPPI.org.uk)

Table: Rate (%) Cancer Screening Rates, 2016

	Kingston	London	England
Breast	71.9	69.2	75.5
Cervical	68.3	66.7	72.7
Bowel	55.3	48.8	57.9

Source: Public Health England, 2018

Table: Percentage of Adults classified as Overweight or Obese in Kingston, London and England, 2016-17

Kingston	London	England
53.4	55.2	61.3
(48.7 to 58.3)	(54.5 to 56.0)	(61.0 to 61.5)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England (based on Active Lives survey, Sport England), 2018

Table: BMI Categories (expressed in percentage) of adults in Kingston, Kingston Lifestyle Survey, 2014

BMI Category	Frequency	Percentage *
Underweight	41	3.00%
Healthy Weight	710	51.20%
Overweight	432	31.10%
Obese	174	12.50%
Severely Obese	30	2.20%
Data Missing/Incorrect entry	178	Not included in percentage
Total people answering	1,565	100.00%

Note: numbers may not add to 100% due to rounding

Source: Kingston Lifestyle Survey, 2014

Table: Modelled estimates of five-a-day consumption of fruit and vegetables in Kingston, London and England, 2006-08

Period	Kingston (%)	London (%)	England (%)
2006-08	40.9	36.4	28.7
	(37.0 to 45.0)	(34.2 to 38.7)	(28.0 to 29.3)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England, 2017

Table: Percentage of adults achieving at least 150 minutes of physical activity per week in accordance with UK Chief Medical Officer (CMO) recommended guidelines on physical activity in Kingston, London and England, 2015

	Kingston	London	England
Inactive adults	20.8	28.1	28.7
	(17.3 to 24.3)	(27.4 to 28.8)	(28.4 to 28.9)
Active adults	66.1	57.8	57.0
	(62.1 to 70.2)	(57.1 to 58.6)	(56.8 to 57.3)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England, 2017

Table: Smoking prevalence of people aged 18 and over in Kingston, London and England, 2012-2015

	Kingston	London	England
2015	13.8%	15.2%	15.5%
	(10.8 to 16.9)	(14.6 to 15.7)	(15.3 to 15.7)

Note: Values in brackets denote 95% confidence intervals

In 2013, several survey modules were removed from the Integrated Household Survey (IHS) to leave a dataset solely based upon the Annual Population Survey (APS). Some differences in survey coverage, imputation and weighting methodology may have resulted in some discontinuity for certain 'core' variables compared to estimates previously provided as part of the IHS, therefore the HIS and APS should not be directly compared.

Source: Annual Population Survey 2012-2015, Public Health England, 2017

Table: Smoking prevalence in Kingston, 2014

	Female	Male	Total	Female (%)	Male (%)	Total (%)
Smoker	100	162	262	12.50%	21.30%	16.70%
Ex-smoker	236	229	465	29.40%	30.10%	29.70%
Never smoked	467	371	838	58.20%	48.70%	53.60%
Total	803	762	1,565	100%	100%	100%

Note: Total number of resident participants = 1565

Source: Kingston Lifestyle Survey, 2015

	Female (%)	Male (%)	Total (%)
Smoker	12.50%	21.30%	16.70%
Ex-smoker	29.40%	30.10%	29.70%
Never smoked	58.20%	48.70%	53.60%
Total	100%	100%	100%

Note: Total number of resident participants = 1565

Source: Kingston Lifestyle Survey, 2015

Table: Rate / 100,000 Admission Episodes for Alcoholic Liver Disease Conditions (2016/17)

Kingston	London	England
411	529	636

Source: Public Health England, 2017

Table: Rate/100,000 Early Deaths (under 75 Years) from Liver Disease (2014-2016)

	Kingston	London	England
Female	9.9	10.5	12.8
Male	19.6	23.4	23.9

Source: Public Health England, 2017

Table: Perceptions of drunk or rowdy behaviour as a problem, 2013

Period		Kingston Town	Malden and Coombe	South of the Borough	Surbiton
Oct-13	Percentage of respondents to 'Your Kingston, Your Say' survey who felt people being drunk or rowdy in public places was an issue in their local area	39%	26%	19%	24%

Source: DJS (November 2013), Your Kingston, Your Say Kingston Resident Survey

Table: Drinking behaviour of adults (persons aged 18+) in Kingston, London and England, 2011-14

	Kingston	London	England
Percentage of adults who abstain from drinking alcohol	14.39 (9.31 to 21.55)	24.35 (23.01 to 25.74)	15.48 (15.09 to 15.88)
Percentage of adults binge drinking on heaviest drinking day	13.67 (8.72 to 20.76)	13.18 (12.09 to 14.34)	16.52 (16.10 to 16.94)
Percentage of adults drinking over 14 units of alcohol a week	29.20 (22.05 to 37.46)	21.57 (20.28 to 22.93)	25.74 (25.26 to 26.22)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England, 2017

Table: Estimates of the prevalence of opiate use and/or crack cocaine use per 1,000 population, amongst People aged 15-64, 2014-2015

	Kingston	England
OCU*	4.77 (3.80 to 7.56)	8.57 (8.49 to 8.86)
Opiate Users	3.88 (3.15 to 6.22)	7.33 (7.3 to 7.6)
Crack Users	2.37 (1.80 to 3.40)	5.21 (5.03 to 5.43)

Note: Values in brackets denote 95% confidence intervals

*"OCU" is defined as a client presenting with opiates and/or crack cocaine as their main, second or third drug recorded at any episode during their latest treatment journey.

Source: National Treatment Agency, Public Health England, 2017

Table: Overall drug treatment activity in 2015-16 compared to 2014-15

	Numbers in treatment		New presentations to treatment	
Opiate	239	No change	85	Down 1%
Non-opiate	50	Up 16%	39	Up 26%
Non-opiate and alcohol	78	Down 16%	60	Down 19%
Total	367	Down 2%	184	Down 4%

Source: Adults - Drugs JSNA Support pack, National Drug Treatment Monitoring System, 2017

Table: Successful completion of drug treatment for opiate and non-opiate users (% who do not re-present within 6 months), aged 18 to 75 years in Kingston, London and England, 2016

	Kingston (%)	London (%)	England (%)
Opiate users	8.2 (5.3 to 12.6)	7.2 7.6	(6.9 to 6.7 (6.6 to 6.8)
Non Opiate users	46.2 (37.5 to 55.2)	38.7 39.7	(37.8 to 37.1 (36.7 to 37.5)

Note: Values in brackets denote 95% confidence intervals

Source: Calculated by Public Health England: Knowledge and Intelligence Team (North West) using data from the National Drug Treatment Monitoring System, 2018

Table: HIV testing uptake and coverage for Kingston, London and England (all Ages), 2016

	Kingston	London	England
HIV testing uptake total (%)	74.52 (73.53 to 75.48)	78.81 78.93	(78.70 to 76.49 (76.42 to 76.56)
HIV testing coverage total (%)	68.51 (67.42 to 69.59)	72.76 72.89	(72.63 to 67.72 (67.65 to 67.80)

Note: Values in brackets denote 95% confidence intervals

Source: Sexual Health Profile, Public Health England, 2017

Table: Percentage of over 25s who choose LARC as their main method of contraception in SRH Services (2016)

Kingston	London	England
30.9	34.4	35.7

Source: Reproductive and Sexual Health Profile, Public Health England, 2018

Table: Total Abortion rate / 1000 (2016)

Kingston	London	England
14.5	20.8	16.7

Source: Reproductive and Sexual Health Profile, Public Health England, 2018

Table: Long-term health problem or disability in Kingston, London and England, persons, all usual residents, Census 2011

Disability	Kingston	London	England	Kingston (%)	London (%)	England (%)
Day-to-day activities limited a lot	8,605	551,664	4,405,394	5.40%	6.70%	8.30%
Day-to-day activities limited a little	11,297	605,501	4,947,192	7.10%	7.40%	9.30%
Day-to-day activities not limited	140,158	7,016,776	43,659,870	87.60%	85.80%	82.40%
Total	160,060	8,173,941	53,012,456	100%	100%	100%

Source: Census 2011, Office for National Statistics, Table QS303EW Accessible via Nomis Web

Table: Cardiovascular Disease Profile of Kingston, similar CCGs, South West London Sustainability and Transformation Plan (STP), and England, 2014-2016

Key Facts	Kingston CCG	Similar CCGs	STP	England
Coronary Heart Disease prevalence (%)	2.1	2.6	2	3.2
Heart Failure prevalence (%)	0.5	0.6	0.5	0.8
Coronary Heart Disease admissions (rate per 100,000)	519.4	-	519	515.8
Heart Failure admissions (rate per 100,000)	171.9	-	168.5	156.9
CHD early mortality (rate per 100,000)	31.7	-	35.3	39.4

Source: CVD Profile: Health Disease, Public Health England, Feb 2018

Table: Recorded prevalence of cardiovascular diseases, Kingston CCG, all ages, 2016-17

	Prevalence (%)
Atrial Fibrillation	1.2
CVD-PP	1.2
CHD	2.1
Heart Failure	0.5
Hypertension	10.3
Peripheral Arterial Disease	0.33
Stroke and TIA	1.01

Note: *CVD Primary Prevention data is for age group 30 to 74 years. Data for years prior to 2014-15 was recorded for all ages and hence is not directly comparable.

Source: Quality and Outcomes Framework (QOF) for April 2015 - March 2016, England

Table: Age-standardised rate of mortality from all cardiovascular diseases per 100,000 population in Kingston, London and England, 2014-16

	Kingston	London	England
Under 75 mortality rate from all cardiovascular diseases	61.7 (53.5 to 70.8)	74.9 76.3	(73.5 to 73.5 (73.0 to 73.9)
Under 75 mortality rate from cardiovascular diseases	38.9 (32. to 46.2)	46.2 (45.1 to 47.3)	46.7 (46.4 to 47.1)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England, 2017

Table: Stroke Statistics for Kingston CCG and England

	Age group	Period	Kingston CCG	London	England
Stroke: QOF prevalence	All ages	2016/17	1.00 (1.00 to 1.10)	1.1 (1.00 to 1.10)	1.70 (1.70 to 1.70)
Stroke mortality rates, DSR per 100,000 population (age standardised)	<75 years	2014-16	12.00 (8.50 to 16.40)	-	13.40 (13.20 to 13.60)
Stroke mortality rates, DSR per 100,000 population (age standardised)	75+ years	2014-16	387.50 (324.9 to 458.5)	-	563.8 (559.8 to 567.80)
Stroke admissions (DSR per 100,000 population)	All ages	2016/17	157	-	7709

Stroke all age admission trends (DSR per 100,000 population)	All ages	2016/17	189.4 (166.00 to 215.10)	-	169.2 (168.1 to 170.3)
Stroke patients who are assessed at 6 months	All ages	2016/17	15.7	-	31.6
Stroke admissions with history of atrial fibrillation not prescribed anticoagulation prior to stroke	All ages	2016/17	33.3	-	47.5

Note: Values in brackets denote 95% confidence intervals
Source: Public Health England, 2017

Table: Estimated Prevalence (%) of Common Mental health Disorders (Population aged 16-74), 2012

	Kingston	London	England
Mixed anxiety and depressive disorder	7.6	8.5	8.9
Generalised anxiety disorder	4.8	5.3	4.5
Depressive episode	3.2	3.5	2.5
All phobias	2	2.2	1.8
Obsessive compulsive disorder	1.4	1.6	1.1
Panic disorder	0.8	0.9	0.7

Source: APMS

Table: Prevalence of Common Mental Health Disorders in Kingston, London and England

	Period	Kingston	London	England
Mental Health: QOF prevalence (all ages)	2015/16	0.84 (0.80 to 0.88)	1.08	0.90 (0.90 to 0.90)
Depression: Recorded prevalence (aged 18+)	2015/16	6.21 (6.09 to 6.33)	6.04 (6.02 to 6.06)	8.26 (8.26 to 8.27)
Depression and anxiety prevalence (GP Patient Survey): % of respondents aged 18+	2015/16	9.91 to 11.1)	(8.84	11.83
Mixed anxiety and depressive disorder: estimated % of population aged 16-74	2012	7.63	8.5	8.92
Generalised anxiety disorder: estimated % of population aged 16-74	2012	4.76	5.34	4.5
Depressive episode: estimated % of population aged 16-74	2012	3.19	3.53	2.48
All phobias: estimated % of population aged 16-74	2012	1.95	2.19	1.77
Obsessive compulsive disorder: estimated % of population aged 16-74	2012	1.41	1.55	1.1
Panic disorder: estimated % of population aged 16-74	2012	0.75	0.86	0.65
Estimated prevalence of post-traumatic stress disorder (PTSD): % of population aged 16+	2012	2.98	3.04	3.02

Note: Values in brackets denote 95% confidence intervals
Source: Public Health England, 2017

Table: Self-reported emotional wellbeing of people aged 16 years and over in Kingston, 2014

	None of the time	Rarely	Some of the time	Often	All of the time	Don't Know
Optimistic about the future	9.33% (6.86 to 11.80)	8.12% (5.65 to 10.59)	28.75% (26.28 to 31.22)	29.78% (27.31 to 32.25)	19.23% (16.76 to 21.70)	4.79% (2.32 to 7.26)
Useful	4.98% (2.51 to 7.45)	5.24% (2.77 to 5.24)	22.43% (19.96 to 22.43)	36.04% (33.57 to 36.04)	27.99% (25.52 to 27.99)	3.32% (0.85 to 3.32)
Relaxed	7.54% (5.07 to 10.01)	14.70% (12.23 to 14.70)	32.97% (30.50 to 32.97)	29.39% (26.92 to 29.39)	13.48% (11.01 to 13.48)	1.92% -
That you are dealing with problems well	4.41% (1.94 to 6.88)	5.50% (3.03 to 7.97)	23.07% (20.60 to 25.54)	41.41% (38.94 to 43.88)	22.75% (20.28 to 25.22)	2.88% (0.41 to 5.35)
That you are thinking clearly	3.51% (1.04 to 5.98)	2.49% (0.02 to 4.96)	15.46% (12.99 to 17.93)	39.94% (37.47 to 42.41)	36.23% (33.76 to 38.70)	2.36% -
That you are able to make up your own mind about things	2.94% (0.47 to 5.41)	2.11% -	9.65% (7.18 to 12.12)	31.88% (29.41 to 34.35)	51.25% (48.78 to 53.72)	2.17% -
That you are close to other people	3.32% (0.85 to 5.79)	2.81% (0.34 to 5.28)	12.01% (9.54 to 14.48)	31.05% (28.58 to 33.52)	48.18% (45.71 to 50.65)	2.62% (0.15 to 5.09)

Note: - indicate that number of responses were too small to allow calculation of the confidence interval.
Number of people who responded to this question was 1,565.
Source: Kingston Lifestyle Survey, 2014

Table: Percentage of Breast, Bowel, and Cervical Cancer Screening (2016/17).

	Kingston	London	England
Persons aged 60-69 screened for bowel cancer within 6 months of invitation (% uptake)	53.4	46.9	57
Females aged 50-70 screened for Breast Cancer within 6 months of invitation (% Uptake)	70.2	65.3	72.2
Females aged 25-64, attending cervical screening within target period (3.5 or 5.5 year coverage, % uptake)	67.4	65.8	72.1

Source: Public Health England, 2017

Table: Percentage One Year Cancer Survival in NHS Kingston CCG and England (2017)

	Kingston (%)	England (%)
Breast	97.3	96.5
Bowel	77.9	77.2
Lung	38	36.8

Source: JSNA pack, Cancer Research UK

Table: QOF prevalence of asthma in Kingston, London and England, 2016-17

	Kingston (%)	London (%)	England (%)
2016-17	4.65	4.62	5.94

Source: Quality and Outcomes Framework, NHS Digital, 2016-17

Table: Age-standardised rate of mortality and those considered preventable from respiratory disease in those aged <75 per 100,000 population in Kingston, London and England, persons, 2014-16

	Kingston	London	England
Under 75 mortality	27.4 (22.0 to 33.7)	30.3 (29.4 to 31.2)	33.8 (33.5 to 34.1)
Under 75 mortality considered preventable	14.3 (10.5 to 19.1)	16.5 (15.8 to 17.2)	18.6 (18.3 to 18.8)

Note: Values in brackets denote 95% confidence intervals

Source: Public Health England, 2017